

ON HOPF ALGEBRAS OF DIMENSION pq

PAVEL ETINGOF AND SHLOMO GELAKI

1. MAIN THEOREM

The classification of Hopf algebras of dimension pq , where p, q are distinct prime numbers, over an algebraically closed field k of characteristic 0 is still unknown. In the previous works [EG], [GW] the problem was solved in the semisimple case; namely, it was shown that any semisimple Hopf algebra of dimension pq is trivial (i.e. isomorphic to either a group algebra or to the dual of a group algebra). In the work [N], Ng completed the classification of Hopf algebras of dimension p^2 , which started in [AS], [M]. In addition to that, the problem was also settled in some specific low dimensions. More precisely, Williams did dimensions 6 and 10 [W], Andruskiewitsch and Natale did dimensions 15, 21 and 35 [AN], and Beattie and Dascalescu did dimensions 14, 55 and 77 [BD].

In this paper we contribute to the classification of Hopf algebras of dimension pq by proving the following main theorem. (We shall work over the field of complex numbers, for convenience.)

Theorem 1.1. *Let $p < q$ be odd primes with $q \leq 2p + 1$. Then any complex Hopf algebra of dimension pq is semisimple and hence trivial.*

In particular, this theorem covers all the odd dimensions listed above.

2. PROOF OF THE MAIN THEOREM

Proposition 2.1. *Let H be a finite dimensional Hopf algebra. Let P denote the projective cover of the trivial representation of H , and N be the number of irreducible representations of H . Then $N \dim(P) \leq \dim(H)$.*

Proof. $H = \bigoplus_{V \in \text{Irr}(H)} \dim(V)P(V)$, where $P(V)$ is the projective cover of V . Now, the module $P(V) \otimes {}^*V$ is projective (cf. [KL], p.441, Corollary 2 or [GMS], Proposition 2.1) and $\text{Hom}(P(V) \otimes {}^*V, \mathbb{C}) = \text{Hom}(P(V), V) \neq 0$. Hence $P(V) \otimes {}^*V$ contains P , and its dimension is at least $\dim(P)$. So $\dim(V) \dim(P(V)) \geq \dim(P)$. Adding over all V , we get $N \dim(P) \leq \dim(H)$, as desired. $\square$

Lemma 2.2. *If $P = \mathbb{C}$ then H is semisimple.*

Proof. If $P = \mathbb{C}$ then $\mathbb{C}$ is projective, and hence any representation $V = V \otimes \mathbb{C}$ of H is projective ([KL], p.441, Corollary 2; [GMS], Proposition 2.1). This implies that $\text{Rep}(H)$ is a semisimple category, so H is semisimple. $\square$

Let H be a **non-semisimple** complex Hopf algebra of dimension pq , $p < q$ are primes.

Lemma 2.3. *Let $D \subset \text{Rep}(H)$ be the subcategory of representations whose composition factors are 1-dimensional. Then D is semisimple.*

Proof. Assume that D is not semisimple. Then D contains a 2-dimensional representation V of H which is a nontrivial extension of a character χ_1 by a character χ_2 . This means, V has a basis v_1, v_2 such that for any $a \in H$, one has $av_1 = \chi_1(a)v_1 + f(a)v_2$, $av_2 = \chi_2(a)v_2$. Since $(ab)v_1 = a(bv_1)$, the element $f \in H^*$ satisfies the equality $f(ab) = f(a)\chi_1(b) + \chi_2(a)f(b)$. Thus, $\Delta(f) = f \otimes \chi_1 + \chi_2 \otimes f$, so f is skew-primitive. Since V is a nonsplit extension, f is nontrivial (i.e. not a multiple of $\chi_1 - \chi_2$). Thus, H^* contains nontrivial skew-primitive elements, which contradicts Proposition 1.8 in [AN]. Thus D is semisimple. $\square$

Suppose from now on that p and q are **odd**.

Theorem 2.4. [N] (i) *The order of S^4 is p .*

(ii) *The distinguished grouplike element either in H or in H^* is nontrivial and has order p .*

So we will assume that H^* has distinguished grouplike element χ of order p . Then χ is a 1-dimensional representation of H , $\chi^p = 1$.

Proposition 2.5. *Let V be an irreducible H -module, which is stable under tensoring on the left with χ . Then $\dim(V)$ is divisible by p .*

Proof. As pointed out in the proof of Lemma 2.1 in [AN], $(\text{End} V)^* \subset H^*$ is a Hopf module for $(\mathbb{C}[G(H^*)], H^*)$. Thus the result follows by the Nichols-Zoeller theorem [NZ]. $\square$

Proposition 2.6. *Suppose that V is an irreducible H -module, which is stable under left multiplication by χ . Then $q > 2p + 1$.*

Proof. Assume that $q \leq 2p + 1$. By Proposition 2.5, the dimension of V is divisible by p . We claim that V is the unique irreducible H -module of dimension $\geq p$, and hence the unique irreducible module stable under left multiplication by χ ; in particular $V^* = {}^*V = V$. Indeed, assume that W is another irreducible H -module of dimension $\geq p$. Then

$$pq = \dim(H) \geq \dim(V)^2 + \dim(W)^2 + \sum_{j=1}^p \dim(\chi^j)^2 = 2p^2 + p.$$

Since $q \leq 2p + 1$, this inequality is an equality, and hence H is semisimple, a contradiction.

Assume that Y, Z are other (hence χ -unstable) simple modules such that $Y \otimes Z$ contains V as a constituent of the Jordan-Hölder series. Then $\dim(Y) > 1$, $\dim(Z) > 1$, and either Y or Z (say, Y) has dimension $> \sqrt{p}$, so

$$pq = \dim(H) \geq \sum_{j=1}^p \dim(\chi^j \otimes Y)^2 + (\dim(V))^2 + \sum_{j=1}^p \dim(\chi^j)^2 > 2p^2 + p,$$

i.e. $q > 2p + 1$, a contradiction. Thus, Y, Z do not exist.

Let C be the category of H -modules which do not contain V as a constituent. As we have seen in the previous paragraph, it is a rigid tensor category, so $C = \text{Rep} H'$, where H' is a quotient of H by a nontrivial Hopf ideal. Since χ^j , $1 \leq j \leq p$, are representations of H' , by the Nichols-Zoeller theorem [NZ], the dimension of H' is a divisor of pq which is divisible by p . Hence the dimension of H' is p , and we have $H' = \mathbb{C}[\mathbb{Z}_p]$. In other words, C is semisimple, with simple objects χ^j , $1 \leq j \leq p$.

Let $P(V)$ be the projective cover of V . If $P(V) \neq V$ then the socle of $P(V)$ is $\chi^{-1} \otimes V = V$ (by [EO], Lemma 2.10). Thus, $\dim(P(V)) \geq 2 \dim(V)$, and

$$pq = \dim(H) \geq \dim(V) \dim(P(V)) + \sum_{j=1}^p \dim(\chi^j) \dim(P(\chi^j)) > 2p^2 + p$$

(the $>$ sign is due to the fact that the projective cover of $\mathbb{C}$ cannot equal $\mathbb{C}$, by Lemma 2.2). So, we have a contradiction. Thus $P(V) = V$.

This means that $P(\mathbb{C})$ must contain V as a constituent (otherwise by Lemma 2.3, one has $P(\mathbb{C}) = \mathbb{C}$, hence by Lemma 2.2 H is semisimple, which is a contradiction). Hence $P(\chi^j)$ contains V as a constituent for all j . Thus the dimension of $P(\chi^j)$ is at least $p + 2$ (it involves χ^j , V , and the socle χ^{j-1}). So

$$pq = \dim(H) \geq \sum_{j=1}^p \dim(P(\chi^j)) + (\dim(V))^2 \geq p(p+2) + p^2 = p(2p+2),$$

a contradiction. $\square$

Assume from now on that $\mathbf{q} \leq 2\mathbf{p} + 1$.

Lemma 2.7. *One has, $\dim(P) < p$.*

Proof. By Proposition 2.6, all simple H -modules are χ -unstable. By Lemma 2.3, not all simple H -modules are 1-dimensional. Thus there are at least 2 orbits of simple H -modules under left multiplication by χ . Consider two cases.

1. The number of orbits is ≥ 3 . Then by Proposition 2.1, $\dim(P) \leq q/3 \leq \frac{2p+1}{3} < p$.

2. The number of orbits is 2. Then by Proposition 2.1, $\dim(P) \leq q/2$, so $\dim(P) \leq p$. So it remains to show that $\dim(P) \neq p$. Suppose $\dim(P) = p$. Let V be an element of the unique orbit of non-1-dimensional representations, and suppose $\dim(V) = d$. Then $P(V) \otimes^* V$ contains P , so $\dim(P(V)) \geq p/d$. Clearly, we cannot have $P = P(V) \otimes^* V$, as the dimension of P is a prime. Thus, $P(V) \otimes^* V = P \oplus Q$, where Q is a nontrivial projective module. Thus, $\dim(Q) \geq p/d$ (as Q contains $P(\chi^j) = \chi^j \otimes P$, or it contains $P(\chi^j \otimes V) = \chi^j \otimes P(V)$, because they are the only indecomposable projective H -modules). So $\dim(P(V)) \dim(V) \geq p + p/d = p(d+1)/d$. This implies (like in the proof of Proposition 2.1) that

$$\begin{aligned} pq &\geq \sum_{j=1}^p \dim(P(\chi^j \otimes V)) \dim(\chi^j \otimes V) + \sum_{j=1}^p \dim(P(\chi^j)) \\ &\geq p^2(d+1)/d + p^2 = p^2 \frac{2d+1}{d} > p(2p+1) \end{aligned}$$

(since clearly $d < p$: otherwise $pq = \dim(H) \geq p \dim(V)^2 = p^3$, which is absurd). This is a contradiction, so the lemma is proved. $\square$

Lemma 2.8. *For any simple H -module V , one has $\chi^{-1} \otimes V \otimes \chi = V$.*

Proof. Let Y, Z be irreducible representations of H which are stable under conjugation by χ , and suppose that $Y \otimes Z$ contains as a constituent an irreducible representation X which is not stable under conjugation by χ . Then $Y \otimes Z$ must also contain as constituents all the translates $\chi^i \otimes X \otimes \chi^{-i}$, which are pairwise non-isomorphic. To be definite, assume that $\dim(Y) \geq \dim(Z)$. Since X cannot

be 1-dimensional, we find that $\dim(Y)\dim(Z) \geq 2p$, so $\dim(Y)^2 > 2p$. This means that

$$pq = \dim(H) \geq \sum_{j=1}^p (\dim(\chi^j \otimes Y)^2 + \dim(\chi^j)^2) > 2p^2 + p,$$

a contradiction. Thus Y, Z do not exist.

So let E be the category of representations of H whose all simple constituents are stable under conjugation by χ . As we have just shown, it is a rigid tensor category. So $E = \text{Rep}(H')$, where H' is a quotient of H .

Assume $H' \neq H$. Then $H' = \mathbb{C}[\mathbb{Z}_p]$. This means that all simple modules over H which are not 1-dimensional fail to be invariant under conjugation by χ .

Choose a non-1-dimensional simple constituent V in P . Such exists, otherwise by Lemma 2.3, $\mathbb{C}$ is projective and H is semisimple, which is a contradiction. Then all translates $\chi^i \otimes V \otimes \chi^{-i}$ (which are pairwise non-isomorphic) are contained in P as constituents. Thus $\dim(P) \geq 2p$ (since $\dim(V) \geq 2$). But we showed before that $\dim(P) < p$. Contradiction. Thus, $H = H'$, as desired. $\square$

Corollary 2.9. *For any simple H -module V , one has $V = V^{****}$.*

Proof. Follows from Lemma 2.8 and the Radford S^4 formula [R]. $\square$

Corollary 2.10. *There exists an odd dimensional indecomposable projective module Q over H such that $Q = Q^{**}$.*

Proof. We have $H = \bigoplus \dim(V)P(V)$ and $H^{**} = H$. By Corollary 2.9, $**$ is an involution on the set of indecomposable projectives $P(V)$. Since the dimension of H is odd, the sum $\bigoplus_{V=V^{**}} \dim(V)P(V)$ is also odd dimensional. Thus one of the summands is odd dimensional, as desired. $\square$

Lemma 2.11. *Let Q be an odd dimensional indecomposable projective module over H such that $Q = Q^{**}$. Then $\dim(Q) \geq p$.*

Proof. Let $a : Q \rightarrow Q^{**}$ be an isomorphism. In other words, a is a linear operator $Q \rightarrow Q$ such that $ax = S^2(x)a$, $x \in H$. We claim that if λ, μ are eigenvalues of a then λ/μ is a root of unity of degree $2p$, i.e. a power of $z = e^{\pi i/p}$.

Indeed, assume that this is not the case. Let Q_1 be the sum of generalized eigenspaces of a on Q with eigenvalues λz^m , $m = 0, \dots, 2p-1$, and Q_2 the sum of generalized eigenspaces of a with other eigenvalues. Then $Q_1, Q_2 \neq 0$, and $Q_1 \oplus Q_2 = Q$. We claim that Q_1 and Q_2 are H -submodules of Q (this is a contradiction, since Q is indecomposable). Indeed, let x be an eigenvector of S^2 in H with eigenvalue z^m , and $v \in Q_i$ for some i be a generalized eigenvector of a with eigenvalue β : $(a - \beta)^N v = 0$. Since $ax = S^2(x)a = z^m xa$, we have $0 = x(a - \beta)^N v = (z^{-m}a - \beta)^N xv$. This shows that xv is a generalized eigenvector of a with eigenvalue βz^m , i.e. $xv \in Q_i$ for the same i . But by Theorem 2.4, elements x as above span H . We are done.

Let us normalize a in such a way that one of its eigenvalues is 1. Then, as we have shown, other eigenvalues will be z^j , $j = 0, \dots, 2p-1$.

By Lorenz' theorem [L], (see also, [EO], proof of Theorem 2.16), $\text{Tr}(a) = 0$. Thus, if m_j is the multiplicity of the eigenvalue z^j , then

$$\sum_{j=0}^{2p-1} m_j z^j = 0, \quad \sum_{j=0}^{2p-1} m_j = \dim(Q).$$

Since $z^{p+j} = -z^j$, we can rewrite the last equation as

$$\sum_{j=0}^{p-1} (m_{2j} - m_{2j+p}) z^{2j} = 0.$$

This means that $m_{2j} - m_{2j+p} = b$ for all j . Since $\sum_j m_j$ is odd, $b \neq 0$. Thus, $\max(m_{2j}, m_{2j+p}) \geq 1$, and hence $\dim(Q) \geq p$, as desired. $\square$

Remark 2.12. The proof of Lemma 2.11 is similar to [N], Section 5.

Lemma 2.13. *The module P is odd dimensional.*

Proof. By Corollary 2.10 and Lemma 2.11, some indecomposable projective module $Q = P(V)$ over H has dimension $\geq p$. If V is not 1-dimensional, then

$$pq = \dim(H) \geq \sum_{j=1}^p \dim(P(\chi^j \otimes V)) \dim(\chi^j \otimes V) + \sum_{j=1}^p \dim(P(\chi^j)) \dim(\chi^j) > 2p^2 + p$$

(since $\dim(V) \geq 2$ and $P(\mathbb{C}) \neq \mathbb{C}$ by Lemma 2.2), a contradiction. So V is 1-dimensional, hence we may assume that $V = \mathbb{C}$, as desired. $\square$

Now we observe that Lemma 2.13 and Lemma 2.11 imply that $\dim P \geq p$, which contradicts our previous conclusion (Lemma 2.7) that $\dim(P) < p$. Thus we get an ultimate contradiction, which means that nonsemisimple Hopf algebras of dimension pq with $p < q \leq 2p + 1$, where p, q are primes, do not exist. The main theorem is proved.

3. ACKNOWLEDGMENTS

The authors thank V. Ostrik for useful discussions. P.E was partially supported by the NSF grant DMS-9988796. P.E partially conducted his research for the Clay Mathematics Institute as a Clay Mathematics Institute Prize Fellow. S.G's research was supported by Technion V.P.R. Fund - Dent Charitable Trust- Non Military Research Fund, and by THE ISRAEL SCIENCE FOUNDATION (grant No. 70/02-1).

REFERENCES

- [AN] N. Andruskiewitsch and S. Natale, Counting arguments for Hopf algebras of low dimension, *Tsukuba J. Math.* **25** (2001), no. 1, 187–201.
- [AS] N. Andruskiewitsch and H-J. Schneider, Hopf algebras of order p^2 and braided Hopf algebras of order p *J. Algebra* **199** (1998), no. 2, 430–454.
- [BD] M. Beattie and S. Dascalescu, Hopf algebras of dimension 14, *preprint*, math.QA/0205243.
- [EG] P. Etingof and S. Gelaki, Semisimple Hopf algebras of dimension pq are trivial, *Journal of Algebra* **210** (1998), 664–669.
- [EO] P. Etingof and V. Ostrik, Finite tensor categories, *preprint*, math.QA/0301027.
- [GMS] E. L. Green, E. N. Marcos and Solberg, Representations and almost split sequences for Hopf algebras, Representation theory of algebras (Cocoyoc, 1994), 237–245, CMS Conf. Proc., **18**, Amer. Math. Soc., Providence, RI, 1996.
- [GW] S. Gelaki and S. Westreich, On semisimple Hopf algebras of dimension pq , *Proceedings of the AMS*, **128** (2000), no.1, 39–47.
- [KL] D. Kazhdan and G. Lusztig, Tensor structures arising from affine Lie algebras. IV. *J. Amer. Math. Soc.* **7** (1994), no. 2, 383–453.
- [L] M. Lorenz, Representations of finite-dimensional Hopf algebras, *J. Alg.* **188** (1997), no. 2, 476–505.

- [M] A. Masuoka, The p^n theorem for semisimple Hopf algebras, *Proc. Amer. Math. Soc.* **124** (1996), no. 3, 735–737.
- [N] S-H. Ng, Non-semisimple Hopf algebras of dimension p^2 *J. Algebra* **255** (2002), no. 1, 182–197.
- [NZ] W.D. Nichols and M.B. Zoeller, A Hopf algebra freeness theorem, *American Journal of Mathematics* **111** (1989), 381–385.
- [R] D.E. Radford, The order of the antipode of a finite-dimensional Hopf algebra is finite, *Amer. J. Math.* **98** (1976), 333–355.
- [W] R. Williams, Finite Dimensional Hopf algebras, *Ph.D thesis, Florida State University*, 1998.

DEPARTMENT OF MATHEMATICS, MASSACHUSETTS INSTITUTE OF TECHNOLOGY, CAMBRIDGE, MA 02139, USA

E-mail address: `etingof@math.mit.edu`

DEPARTMENT OF MATHEMATICS, TECHNION-ISRAEL INSTITUTE OF TECHNOLOGY, HAIFA 32000, ISRAEL

E-mail address: `gelaki@math.technion.ac.il`